

Anti-p53 Phospho-Ser20 antibody, mouse monoclonal (17B6)

Product code	71-113
Size	100 µg
Storage	-20°C
Concentration	1.0 mg/ml
Buffer	PBS- with 50% glycerol
Purity	Purified IgG fraction with protein A from hybridoma cell culture medium.
Immunogen	Synthetic peptide containing phosph-Ser20
Isotype	Mouse IgG1 κ
Reactivity	Human p53 phosphorylated at Ser20
Validation	N/A
Application	1. Western blotting (~1/1,000 dilution) 2. ELISA
Background	<p><i>p53</i> mutants are found in more than half of human cancers and are considered as the most important human cancer related gene. <i>p53</i> is detected at 53kD position by electrophoresis and is composed of 393 amino acids. In the unstressed normal cells, the <i>p53</i> level is low and it is inactive. However, with stress, especially with DNA damage, it is activated to promote arrest of cell cycle and repair of DNA damage, or induction of apoptosis. The functions of <i>p53</i> are regulated by phosphorylation of serine and threonine, and acetylation of lysine at various sites in the molecule (1).</p> <p><i>p53</i> at Ser20 is phosphorylated by Chk1 and Chk2 kinases and this enhances tetramerization, stability and activity of <i>p53</i> (2, 3).</p>
Data Link	UniProtKB/Swiss-Prot P04637 (P53_HUMAN)
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

Data Images: 71-113 Anti-p53 Phospho-Ser20 antibody, mouse monoclonal (17B6)




Fig.1 Identification of p53 phospho-Ser20 with monoclonal antibody 17B6 by Western blotting.

The lysates of H1299 cells (p53 non-producing) transfected either with Flag vector (left lane), Flag-p53 (wild-type) expressing plasmid (middle lane) or Flag-p53 mutant (Ser20Ala) expressing plasmid (right lane) were immunoprecipitated with anti-Flag antibody and then analyzed by Western blotting with anti-p53 p-S20 monoclonal antibody, 17B6, (upper panel) or with pantropic p-53 antibody (lower panel).

Related Products: Many antibodies specific to phosphorylated and acetylated oncogene products

#71-115 anti-p53 (p-S46)

#71-117 anti-p53 (p-S315)

#71-131 anti-p53 (Ac-K120)

#71-133 anti-p53 (Ac-K382)

References

1. Bode AM & Dong Z "Post-translational modification of p53 in tumorigenesis" *Nature Rev Cancer* **4**: 793 -805 (2004) PMID: [15510160](#)
2. Shieh SY *et al* "DNA damage-inducible phosphorylation of p53 at N-terminal sites including a novel site, Ser20, requires tetramerization" *EMBO J* **18**: 1815-1823 (1999) PMID: [10202145](#)
3. Hirao A "DNA damage-induced activation of p53 by the checkpoint kinase Chk2" *Science* **287**: 1824-1827 (2000) PMID: [10710310](#)