

Anti-Brg1 antibody, rabbit polyclonal, affinity-purified

Product code	70-230
Size	100 µg
Storage	-20°C
Concentration	1.0 mg/ml
Buffer	PBS- with 50% glycerol
Purity	Affinity-purified with immunogen peptide from rabbit antiserum
Immunogen	C-terminal 50 amino acids of human BRG1, C-KLGRKEKAQDRLK GGRRRPSR GSRAKPVVSDDDSEEEQEEDRSRGSGSEED, conjugated with KLH
Isotype	IgG
Reactivity	Human and mouse. Not tested in other species
Special notes	N/A
Application	<ol style="list-style-type: none"> 1. Western blotting (1/1,000 dilution) (Fig. 1) 2. Immunoprecipitation (Assay dependent) 3. Immunofluorescence staining (1/1,000 dilution) (Fig. 3) <p>Not tested for other applications.</p>
Background	<p>Brg1 (1647 aa, 185 kDa) is transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. Component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator complex. In resting neurons, transcription of the c-FOS promoter is inhibited by BRG1-dependent recruitment of a phospho-RB1-HDAC repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex. At the same time, there is increased recruitment of CREBBP to the promoter by a CREST-dependent mechanism, which leads to transcriptional activation. The CREST-BRG1 complex also binds to the NR2B promoter, and activity-dependent induction of NR2B expression involves a release of HDAC1 and recruitment of CREBBP. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth. SMARCA4/BAF190A may promote neural stem cell self-renewal/proliferation by enhancing Notch-dependent proliferative signals, while concurrently making the neural stem cell insensitive to SHH-dependent differentiating cues By similarity. Also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. Acts as a corepressor of ZEB1 to regulate E-cadherin transcription and is required for induction of epithelial-mesenchymal transition (EMT) by ZEB1.</p>
Data Link	UniProtKB P51532 (SMCA4_HUMAN)
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.	

Data Images: 70-230 Anti-Brg1 antibody, rabbit polyclonal, affinity-purified




Fig.1 Identification of Brg1 in whole cell extracts of human and mouse cells by western blotting using anti-Brg1 antibody.

Lane 1; HeLa (human) cell extract

Lane 2; HeLa (human) treated Adriamycin cell extract

Lane 3; NIH 3T3 (mouse) cell extract

*Star indicates the position of Brg1 protein bands (Predicted molecular mass of Brg1 is 185 kDa).

Anti-Brg1 antibody was used at 1/1,000 dilution




Fig 2. Immunoprecipitation of Brg1 protein from nuclear extracts of HeLa cells by using anti-Brg1 antibody.

Brg1 was precipitated with anti-Brg1 antibody and proteinG-conjugated agarose beads and probed with anti-Brg1 antibody by western blotting.

Lane1; Input nuclear extracts

Lane 2; Control IP with non-immune IgG

Lane 3; Immunoprecipitation with anti-Brg1 antibody




Fig.3. Immunofluorescence staining of Brg1 protein in HeLa cells.

HeLa cells were fixed in 4% paraformaldehyde overnight and permeabilized in 0.25% TritonX 100 in PBS for 10 min. Anti-Brg1 antibody was used at 1/1,000 dilution. As second antibody, goat anti-rabbit IgG conjugated with Alex488 was used at 1/1,000 dilution. Nuclei were stained with DAPI. Brg1 protein is localized in nuclei.




Fig 4. Silencing of Brg1 expression by Brg1-specific siRNA

HeLa cells were transfected with the siRNA described in the paper. 48 hr later, Brg1 in total cell extracts was detected by Western blotting. The signal intensity of the blot was scanned and displayed as 100% of control sample.

Publication: This antibody has been described and used in the following publication.

Nishimoto N. et al (2012) Heterocomplex formation by Arp4 and β -actin is involved in the integrity of the Brg1 chromatin remodeling complex. J Cell Sci.125: 3870-82. [PubMed](#) [WB](#), [IP](#)