

## Anti-RPC19 (*S. cerevisiae*) antibody, rabbit serum

62-035 100  $\mu$ l

**Shipping and Storage:** Shipped at 4°C or -20°C and store at -20°C

**Immunogen:** Recombinant GST-RPC19 fusion protein

**Form:** Whole antiserum added with 0.09 % sodium azide

**Reactivity:** *S. cerevisiae* RPC protein. Not tested in other species.

### Applications

- 1) Western blotting (1/1,000) dilution)
- 2) ELISA

**Background:** DNA-dependent RNA polymerases catalyze the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Common core component of RNA polymerases I and III which synthesize ribosomal RNA precursors and small RNAs, such as 5S rRNA and tRNAs, respectively.

From the sequence data, RPC19 consists of 142 amino acids with molecular mass of 16,151 Da.

**Data Link:** UniProt [P28000](#) DNA-directed RNA polymerases I and III subunit RPAC2

**References:** This antibody was described and used in the following publication.

Todaka Y. et al. Association of the GTP-Binding Protein Gtr1p with Rpc19p, a Shared Subunit of RNA Polymerase I and III in yeast *Saccharomyces cerevisiae*. [Genetics](#). 2005 Aug;170(4):1515-24. Epub 2005 Jun 3.

