

Anti- *V. parahemolyticus* TRH Toxin antibody, rabbit IgG

64-015 100 µg

Shipping and Storage: Shipped at 4°C or -20°C. Store at -20°C.

Immunogen: TRH toxoid and toxin purified from culture medium of *V. parahemolyticus* TRH⁺ strain.

Form: IgG fraction purified from the rabbit anti-TRH serum by salting out and ion-exchange chromatography. 1 mg/ml in PBS- with 50% glycerol. Filter-sterilized. Azide- and carrier-free.

Reactivity: TRH toxin and TDH toxin of *V. parahemolyticus* (Fig. 3)

Applications

- 1) Western blotting (2,000~10,000 dilution) (Fig. 1)
- 2) Immunoprecipitation (Fig. 2)
- 3) ELISA (assay dependent)

Background: Many *Vibrio parahemolyticus* isolated as a cause of food poisoning, produce toxin called hemolysin, and this is the main cause of illness. The hemolysin of *V. parahemolyticus* mainly interacts with an intestinal tract or the heart, produces diarrhea by enterotoxicity, and also there are severe cases of making a patient die by cardiotoxicity. Two kinds, heat-resistant hemolysin (TDH, thermostable direct hemolysin) and heat-resistant toxin related hemolysin (TRH, TDH-related hemolysin), are known as hemolysins of *V. parahemolyticus*. Among these, TDH is known for many years and has been studied more extensively. In order to distinguish whether it is the *V. parahemolyticus* that produces TDH, samples are grown on the Agatsuma medium (blood agar which is added with mannitol), and judged by whether a hemolysis is observed or not. This hemolysis was one of the examining methods, which is called Kanagawa phenomenon to judge whether it is pathogenic *V. parahemolytica*. However, it became clear that the food poisoning by the *V. parahemolyticus* of Kanagawa-phenomenon negativity was discovered, and this organism did not produce TDH, but it produced TRH. Moreover, since the Kanagawa phenomenon sensitivity is not so high, the immunological technique employing antibody against toxin is used together for the judgment of pathogenicity of *V. parahemolyticus*.

TRH is the heat labile toxin protein of molecular weight 21.1 kDa (189 aa). Homology of TDH (21.3 kDa, 189 aa) with TRH is about 60% (reference 1 and 2), and shows partial antigenic similarities. Susceptibility of the blood cells of various animals to TRH differs greatly, and TRH shows more than 100 times in rabbit skin capillary permeability activity than TDH.

Data linkage

[GenBank BAB13778.1](#) TDH-related hemolysin [*Vibrio parahaemolyticus*]

[UniProKB P19249](#) Thermostable direct hemolysin 1

References

1. Honda T. Et al . (1988) Purification and characterization of a hemolysin produced by a clinical isolate of Kanagawa phenomenon-negative *Vibrio parahaemolyticus* and related to the thermostable direct hemolysin. *Infect. Immun.* 56 :961-965. . [PubMed ID 3126151](#) **Free access**
2. Nishibuchi M et al. (1989) Cloning and nucleotide sequence of the gene (trh) encoding the hemolysin related to the thermostable direct hemolysin of *Vibrio parahaemolyticus* . *Infect Immun* 57:2691-7. [PubMed ID 2759706](#) **Free access**






Fig. 1. Detection of TRH toxin in the culture medium of *V. parahaemolyticus* (TRH⁺) by western blotting with anti-TRH antibody.

1. Ten-times condensed sample were analyzed by SDS-PAGE and stained with CBB.
2. The same sample was analyzed by western blotting. The antibody at 1/2,000 dilution was used. TRH toxin band is indicated by an arrow.

Fig. 2. Immunoprecipitation of TRH toxin from *Vibrio parahaemolyticus* (TRH⁺) culture medium with anti-TRH antibody.

1. TRH was precipitated with anti-TRH antibody from the same sample as in Fig.1, analysed by SDS-PAGE and stained with CBB.
2. TRH was precipitated with anti-TRH antibody from the same sample and TRH was detected by WB with anti-TRH antibody.

Arrow indicates TRH band. H and L indicate heavy and light chain of IgG, respectively.

Fig.3. Anti-TRH antibody cross-reacts with TDH toxin in western blot.

1. Partially purified TDH toxin
 2. Condensed culture medium of *V. haemolyticus*.
- Antibody was used at 1/2,000 dilution for WB.