

Anti-Cnd2 (*S. pombe*) antibody, rabbit serum

63-105 100 µl

Shipping and Storage: shipped at 4°C or -20°C and store at -20°C

Immunogen: Recombinant GST-Cnd2 fusion protein expressed in *E. coli*

Form: Rabbit antiserum added with 0.05 % sodium azide

Reactivity: *S. pombe* Cnd2 protein. Not tested for other species

Applications

1. Western blotting (500 fold dilution) (Ref.1,2,3)
2. Immunofluorescent staining (Ref. 3)

Not tested for other applications

Background: Cnd2 protein is a regulatory subunit of the condensin complex, which is required for conversion of interphase chromatin into mitotic condensed chromosomes in *S. pombe*. The condensin complex also plays a role during interphase in processes of DNA repair and recombination.

Althogh molecular mass of the Cdn2 protein of *S. pombe* from the DNA sequence is 83.2 kDa, it migrates on SDS-PAGE at a position equivalent to a 105 kDa protein (Ref. 2).

Data Link: [cnd2: pombe : GeneDB](#)

References: This antibody was used in the following references.

1. Chen ES, Sutani T, Yanagida M. Chen ES, Sutani T, Yanagida M. (2004) Cti1/C1D interacts with condensin SMC hinge and supports the DNA repair function of condensin. Proc Natl Acad Sci U S A. 101:8078-83. [PNAS Content](#)
2. Aono N, Sutani T, Tomonaga T, Mochida S, Yanagida M. (2002) Cnd2 has dual roles in mitotic condensation and interphase. Nature. 417:197-202. [pubmed/12000964](#)
3. Morishita J.(2001) Bir1/Cut17 moving from chromosome to spindle upon the loss of cohesion is required for condensation, spindle elongation and repair. Genes Cells. 6:743-63. [Genes to Cells - Wiley Online Library](#)




Fig. 1 Identification of Cnd2 protein in the crude extract of *S. pombe* by western blotting with this antiserum.

Fig.2 Detection of Cnd2 protein by immunofluorescent staining of *S. pombe* cell.




Fig.3 Identification of Cnd2 protein and
Cnd2-GFP protein in crude cell extract by
anti-Cnd2 antibody at 1/500 dilution.