

Anti-Nob1 (*S.cerevisiae*) antibody, affinity purified

62-211 100 ul

Background: The 26 S proteasome is a protein complex with a molecular mass of \approx 2,000 kDa. It is essential not only for eliminating damaged or misfolded proteins but also for degrading short lived regulatory proteins involved in cell cycle regulation, DNA repair, signal transduction, apoptosis, and metabolic regulation (1). **Nob1p** is essential nuclear protein required for biogenesis of the 26S proteasome (2). **Nob1p** is speculated to serve as a chaperone to join the 20S proteasome with the 19S regulatory particle in the nucleus and to be degraded upon the maturation of the 26S proteasome (3). **Nob1p** is composed of 459 amino acid residues.

Applications

- 1) Western blotting (~400 fold dilution)
- 2) Immunoprecipitation
- 3) Not tested for other applications

Specification

Product: Rabbit polyclonal antibody affinity purified with recombinant Nob1p

Immunogen: Recombinant yeast Nob1p expressed in E. coli

Form: Purified IgG in PBS, 1 mg/ml BSA, 0.09 % sodium azide, 50% glycerol


Reactivity: *S. cerevisiae* Nob1p. Not tested with other species

Storage: -20°C. Avoid repeated freezing and thawing, store the antibody in aliquots.

Reference: This product was used in ref. 2 and 3

1. Hershko, A., and Ciechanover, A. THE UBIQUITIN SYSTEM *Annu. Rev. Biochem.* **67**, 425-479 (1998)
2. Tone, Y. *et al.* Nob1p, a new essential protein, associates with the 26S proteasome of growing *Saccharomyces cerevisiae* cells. *Gene* **243**:37-45 (2000)
3. Tone, Y., and Toh-e, A. Nob1p is required for biogenesis of the 26S proteasome and degraded upon its maturation in *Saccharomyces cerevisiae*. *Genes & Dev.* **16** :3142-3157 (2002)

Figure1. Detection of Nob1p (51.7 kD) in the crude extract of *S. cerevisiae* by Western blotting using this antibody.



Related products: # 62-201 anti-Rpn3, #62-203 anti-Rpn5,
#62-205 anti-Rpn7, #62-207 anti-Rpn9, #62-209 anti-Rpn12,
#62-213 anti-Nas6, #62-215 anti-Tem1