

Anti-HIV-1 Gag p15 antibody, rabbit serum

Product code	65-011
Size	100 μl
Storage	Store 4°C for short term For long term storage store at -20°C.
	Aliquot to avoid repeated freezing and thawing.
Concentration	N/A
Buffer	0.09% sodium azide
Purity	Rabbit antiserum
Immunogen	Purified full-size recombinant Gag p15 of HIV-1 subtype B (Ref 2) expressed
	in <i>E. coli</i> (Ref 2,3)
Isotype	Rabbit IgG
Reactivity	HIV-1 LAI strain
Special notes	N/A
Application	1. Western blot (1/1,000~1/5000)
11pp110a01011	2. Dot blot (assay dependent)
	3. Immunoprecipitation (assay dependent)
	4. ELISA (assay dependent)
	Other applications have not been tested.
Background	HIV-1 Gag p15 is processed by digestion of its precursor Gag p55 by HIV-1 protease. This protein is further digested into nuclocapsid protein p7 and into p6 and p1 of unknown function. This digestion is promoted by the binding of HIV-1 genome RNA and the two Zn finger motifs that exist in the p7 region. The produced nucleocapsid protein p7 regulates the RNA function by directly binding to HIV-1 genome RNA (1).
Data Link	GenBank: <u>AAA44988.1</u>
Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC	
PROCEDURES. NOT FOR MILITARY USE.	



Data Images: 65-011 Anti-HIV-1 Gag p15 antibody, rabbit serum




Fig.1 Detection of HIV-1 p15 and its p55 precursor protein by Western blotting using the anti-p15 antibody.

Lane1: Extract of MT4 cells

Lane2: Extract of MT4 cells infected with HIV-1 (LAI strain)

The antiserum was diluted 1,000 fold before use.

References

- 1. Freed EO "HIV-1 gag proteins: diverse functions in the virus life cycle" *Virology* **251**:1-15 (1998) Review PMID: 9813197
- 2. Saito A *et al* "Overproduction, purification, and diagnostic use of the recombinant HIV-1 Gag proteins, the precursor protein p55 and the processed products p17, p24, and p15" *Microbiol Immunol* **39**:473-483 (1995) PMID: <u>8569532</u>

Related products

05-007 HIV-1 Gag p15