

## Anti-SCYL2 / CVAK104 antibody, rabbit polyclonal, siRNA validated

71-610 50 µg

**Storage :** Shipped at 4°C or -20°C and store at -20°C

**Validation:** Specificity of reaction was validated with siRNA

**Reactivity:** Human, mouse, rat and hamster.

### Applications

- 1) Western blotting (1/1,000 dilution)
- 2) Immunoprecipitation (1/200-1/1,000 dilution)
- 3) Immunofluorescence staining (1/200-1/1,000 dilution)

**Immunogen:** Human SCYL2 protein (amino acids 528– 929) fused with a His6 tag

**Purity:** Purified IgG (Salting-out and ion-exchange chromatography)


**Form:** 1.0 mg/ml in 1 x PBS and 50% glycerol

**Background:** Component of AP2-containing clathrin coated structures at the plasma membrane or of endocytic coated vesicles. According to PubMed:[15809293](#), probable serine/threonine-protein kinase that phosphorylates, in vitro, the beta2-subunit of the plasma membrane adapter complex AP2 and other proteins in presence of poly-L-lysine. According to PubMed:[16914521](#), has no detectable kinase activity in vitro. May regulate clathrin-dependent trafficking between the TGN and/or the endosomal system

**Data Link:** UniProtKB [Q6P3W7](#) (SCYL2\_HUMAN)

**Reference:** This protein was described and used in the following publication.

Terabayashi T. et al. A coated vesicle-associated kinase of 104 kDa (CVAK104) induces lysosomal degradation of frizzled 5 (Fzd5). [J Biol Chem.](#) (2009) 284(39):26716-24. **WB, IP**




**Fig.1 Validation of the anti-SCYL2 antibody with siRNA.in western blotting.**

293 cells were treated with control of SCYL2-siRNA. At 48 h after transfection,the lysates were analyzed by western blotting with anti-SCYL2 antibody or anti-  $\beta$ -tubulin antibody, the latter for a loading control.

Lane 1: Control siRNA


Lane 2: SCYL2-siRNA



**Fig.2. Detection of endogenous levels of SCYL2 in human and mouse cell extracts by western blotting.**

20  $\mu$ g of lysates of HeLa, MCF7 and NIH3T3 cells were used for western blotting. 7.5% gel was used and blotted overnight in a wet system.

The anti-SCYL2 antibody was used at 1/1,000 dilution and as the 2<sup>nd</sup> antibody, goat anti-rabbit IgG (Abcam 97051) was used at 1/10,000 dilution.



**Fig,3 Immunofluorescence staining of SCYL2 in MCF7 cells.**

MCF7 cells were fixed with 4% PFA and permeabilized with 0.25% Triton X-100 in PBS. The anti-SCYL2 antibody was used at 1/1,000 dilution and as a 2<sup>nd</sup> antibody, goat anti-rabbit IgG conjugated with Alexa Fluor 488 was used at 1/1,000 dilution (left panel). DNA was stained with DAPI (1  $\mu$ g/ml) in TBS. The merged image was shown in the right panel.